

Quasinormal modes and greybody factors of a four-dimensional Lifshitz black hole with $z = 0$

Marcela Catalán* and Eduardo Cisternas†

*Departamento de Ciencias Físicas, Facultad de Ingeniería y Ciencias,
Universidad de La Frontera, Avenida Francisco Salazar 01145, Casilla 54-D, Temuco, Chile.*

P. A. González‡

*Facultad de Ingeniería, Universidad Diego Portales,
Avenida Ejército Libertador 441, Casilla 298-V, Santiago, Chile.*

Yerko Vásquez§

*Departamento de Física, Facultad de Ciencias,
Universidad de La Serena,
Avenida Cisternas 1200, La Serena, Chile.*

(Dated: October 14, 2015)

We study scalar perturbations for a four-dimensional asymptotically Lifshitz black hole in conformal gravity with dynamical exponent $z = 0$, and spherical topology for the transverse section, and we find analytically and numerically the quasinormal modes for scalar fields for some special cases. Then, we study the stability of these black holes under scalar field perturbations and the greybody factors.

PACS numbers:

I. INTRODUCTION

Lifshitz spacetimes have received considerable attention from the condensed matter point of view due to the AdS/CFT correspondence, i.e., searching for gravity duals of Lifshitz fixed points for condensed matter physics and quantum chromodynamics [1]. From the quantum field theory point of view, there are many invariant scale theories of interest when studying such critical points. Such theories exhibit the anisotropic scale invariance $t \rightarrow \chi^z t$, $x \rightarrow \chi x$, with $z \neq 1$, where z is the relative scale dimension of time and space, and these are of particular interest in studies of critical exponent theory and phase transitions. Systems with such behavior appear, for instance, in the description of strongly correlated electrons. The importance of possessing a tool to study strongly correlated condensed matter systems is beyond question, and consequently much attention has focused on this area in recent years.

One of the most well studied systems in the context of gauge/gravity duality, is the holographic superconductor. In its simplest form, the gravity sector is a gravitating system with a cosmological constant, a gauge field and a charged scalar field with a potential. The dynamics of the system defines a critical temperature above which the system finds itself in its normal phase and the scalar field does not have any dynamics. Below the critical temperature the system undergoes a phase transition to a new configuration. From the gravity side this is interpreted as the black hole to acquire hair while from boundary conformal field theory side this is interpreted as a condensation of the scalar field and the system enters a superconducting phase. In this sense, Lifshitz holographic superconductivity has been a topic of numerous studies and interesting properties are found when one generalizes the gauge/gravity duality to non-relativistic situations [2–11].

The Lifshitz spacetimes are described by the metrics

$$ds^2 = -\frac{r^{2z}}{\ell^{2z}} dt^2 + \frac{\ell^2}{r^2} dr^2 + \frac{r^2}{\ell^2} d\vec{x}^2, \quad (1)$$

where \vec{x} represents a $D - 2$ dimensional spatial vector, D is the spacetime dimension and ℓ denotes the length scale in the geometry. If $z = 1$, the spacetime is the usual anti-de Sitter metric in Poincaré coordinates. Furthermore, all

*Electronic address: marceicha@gmail.com

†Electronic address: eduardo.cisternas@ufrontera.cl

‡Electronic address: pablo.gonzalez@udp.cl

§Electronic address: yvasquez@userena.cl

scalar curvature invariants are constant and these spacetimes have a null curvature singularity at $r \rightarrow 0$ for $z \neq 1$, which can be seen by computing the tidal forces between infalling particles. This singularity is reached in finite proper time by infalling observers, so the spacetime is geodesically incomplete [12]. The metrics of Lifshitz black holes asymptotically have the form (1); however, obtaining analytical solutions does not seem to be a trivial task, and therefore constructing finite temperature gravity duals requires the introduction of strange matter content with a theoretical motivation that is not clear. Another way of finding such a Lifshitz black hole solution is by considering carefully-tuned higher-curvature modifications to the Hilbert-Einstein action, as in new massive gravity (NMG) in 3-dimensions or R^2 corrections to general relativity. This has been done, for instance, in [18–21]. A 4-dimensional topological black hole with $z = 2$ was found in [22, 23] and a set of analytic Lifshitz black holes in higher dimensions for arbitrary z in [24]. Lifshitz black holes with arbitrary dynamical exponent in Horndeski theory were found in [25] and non-linearly charged Lifshitz black holes for any exponent $z > 1$ in [26]. Thermodynamically, it is difficult to compute conserved quantities for Lifshitz black holes; however, progress was made on the computation of mass and related thermodynamic quantities by using the ADT method [13, 14] as well as the Euclidean action approach [15, 16]. Also, phase transitions between Lifshitz black holes and other configurations with different asymptotes have been studied in [17]. However, due to their different asymptotes these phases transitions do not occur.

Conformal gravity is a four-derivative theory and is perturbatively renormalizable [27, 28]. Also, it contains ghost-like modes in the form of massive spin-2 excitations. However, a solution to the ghost problem in fourth order derivative theories was shown in [29] by using the method of Dirac constraints [30] to quantize the Pais-Uhlenbeck fourth order oscillator model [31]. In this work, we consider a matter distribution outside the event horizon of the Lifshitz black hole in 4-dimensions in conformal gravity with a spherical transverse section and dynamical exponent $z = 0$. It is worth mentioning that for $z = 0$ the previously mentioned anisotropic scale invariance corresponds to space-like scale invariance with no transformation of time. The matter is parameterized by scalar fields minimally and conformally coupled to gravity. Then, we obtain analytically and numerically the quasinormal frequencies (QNFs) [32–37] for scalar fields, after which we study their stability under scalar perturbations. Also, we compute the reflection and transmission coefficients and the absorption cross section.

The study of the QNFs gives information about the stability of black holes under matter fields that evolve perturbatively in their exterior region, without backreacting on the metric. In general, the oscillation frequencies are complex, where the real part represents the oscillation frequency and the imaginary part describes the rate at which this oscillation is damped, with the stability of the black hole being guaranteed if the imaginary part is negative. The QNFs are independent of the initial conditions and depend only on the parameters of the black hole (mass, charge and angular momentum) and the fundamental constants (Newton constant and cosmological constant) that describe a black hole, just like the parameters that define the test field. On the other hand, the QNFs determine how fast a thermal state in the boundary theory will reach thermal equilibrium according to the AdS/CFT correspondence [38], where the relaxation time of a thermal state is proportional to the inverse of the imaginary part of the QNFs of the dual gravity background, which was established due to the QNFs of the black hole being related to the poles of the retarded correlation function of the corresponding perturbations of the dual conformal field theory [39]. Fermions on a Lifshitz background were studied in [40] by using the fermionic Green’s function in 4-dimensional Lifshitz spacetime with $z = 2$; the authors considered a non-relativistic (mixed) boundary condition for fermions and showed that the spectrum has a flat band. Also, the Dirac quasinormal modes (QNMs) for a 4-dimensional Lifshitz black hole were studied in [41]. Generally, the Lifshitz black holes are stable under scalar perturbations, and the QNFs show the absence of a real part [16, 42–46]. The QNFs have been calculated by means of numerical and analytical techniques, some remarkably numerical methods are: the Mashhoon method, Chandrasekhar-Detweiler, WKB method, Frobenius method, method of continued fractions, Nollert, asymptotic iteration method (AIM) and improved AIM among others. In the context of black hole thermodynamics, QNMs allow the quantum area spectrum of the black hole horizon to be studied [42] as well as the mass and the entropy spectrum.

On the other hand, knowledge of black holes perturbations is also useful for studying the Hawking radiation, which is a semiclassical effect and gives the thermal radiation emitted by a black hole. At the event horizon, the Hawking radiation is in fact blackbody radiation. However, this radiation still has to traverse a non-trivial curved spacetime geometry before reaching a distant observer that can detect it. The surrounding spacetime thus works as a potential barrier for the radiation, giving a deviation from the blackbody radiation spectrum, seen by an asymptotic observer [47]. Thus the total flux observed at infinity is that of a D -dimensional greybody at the Hawking temperature. The factors that modify the spectrum emitted by a black hole are known as greybody factors and can be obtained through the classical scattering (for a review see [48]). In this sense, the scalar greybody factors for an asymptotically Lifshitz black hole were studied in [44, 49], and particle motion on these geometries in [50–52].

The paper is organized as follows. In Sec. II we give a brief review of the 4-dimensional Lifshitz black hole in conformal gravity. In Sec. III we calculate the QNFs of scalar perturbations for the 4-dimensional Lifshitz black hole with spherical topology and $z = 0$ for some special cases analytically and numerically by using the improved AIM. Then, in Sec. IV, we study the reflection and transmission coefficients and the absorption cross section. Finally, our

conclusions are in Sec. V.

II. 4-DIMENSIONAL ASYMPTOTICALLY LIFSHITZ BLACK HOLE IN CONFORMAL GRAVITY

In this work we consider a matter distribution described by a scalar field outside the event horizon of a four-dimensional asymptotically Lifshitz black hole in conformal gravity with $z = 0$ and spherical topology for the transverse section [53]. Conformal gravity is a limit case of Einstein-Weyl gravity. The action of Einstein-Weyl gravity is given by

$$S = \frac{1}{2k^2} \int \sqrt{-g} d^4x \left(\mathcal{R} - 2\Lambda + \frac{1}{2}\alpha |Weyl|^2 \right), \quad (2)$$

where

$$|Weyl|^2 = \mathcal{R}^{\mu\nu\rho\sigma} \mathcal{R}_{\mu\nu\rho\sigma} - 2\mathcal{R}^{\mu\nu} \mathcal{R}_{\mu\nu} + \frac{1}{3}\mathcal{R}^2, \quad (3)$$

\mathcal{R} is the Ricci scalar and Λ is the cosmological constant. When α goes to infinity we have the special case of conformal gravity, and the field equations in vacuum are given by $B_{\mu\nu} = 0$, where $B_{\mu\nu}$ is the Bach tensor defined by:

$$B_{\mu\nu} = (\nabla^\rho \nabla^\sigma + \frac{1}{2}\mathcal{R}^{\rho\sigma}) C_{\mu\nu\rho\sigma}, \quad (4)$$

where $C_{\mu\nu\rho\sigma}$ is the Weyl tensor. The following metric solves the field equations [53]

$$ds^2 = -f dt^2 + \frac{4\ell^2 dr^2}{r^2 f} + r^2 d\Omega_{2,k}^2, \quad (5)$$

$$f = 1 + \frac{\lambda}{r^2} + \frac{\lambda^2 - k^2 \ell^4}{3r^4}. \quad (6)$$

For $k = \pm 1$, there is an event horizon at the largest root of f , given by

$$r_+^2 = \frac{1}{6} \left(\sqrt{3(4\ell^4 - \lambda^2)} - 3\lambda \right), \quad (7)$$

and for $k = 0$ the singularity is naked. Note that the requirement $r_+^2 > 0$ implies that $-2\ell^2 \leq \lambda < \ell^2$. When $\lambda = -2\ell^2$ the solution becomes extremal, and for $k = 1$ the entropy vanishes in this case. The Kretschmann scalar (for $k = 1$) is given by

$$\mathcal{R}_{\mu\nu\rho\sigma} \mathcal{R}^{\mu\nu\rho\sigma} = \frac{9r^8 + 6(\lambda - 4\ell^2)r^6 + (50\ell^4 + \lambda(19\lambda - 24\ell^2))r^4 + 2(\lambda^2 - \ell^4)(21\lambda - 4\ell^2)r^2 + 25(\lambda^2 - \ell^4)^2}{12\ell^4 r^8}, \quad (8)$$

therefore, there is a curvature singularity at $r = 0$. In the next section, we determine the QNFs by considering the Klein-Gordon equation in this background and by establishing the boundary conditions on the scalar field at the horizon and at spatial infinity.

III. QUASINORMAL MODES OF A 4-DIMENSIONAL LIFSHITZ BLACK HOLE

The QNMs of scalar perturbations in the background of a four-dimensional asymptotically Lifshitz black hole in conformal gravity with dynamical exponent $z = 0$ are given by the scalar field solution of Klein-Gordon equation with suitable boundary conditions. This means there are only ingoing waves on the event horizon and we consider that the scalar field vanishes at spatial infinity, known as Dirichlet boundary conditions. These fields are considered as mere test fields, without backreaction over the spacetime itself. Therefore, it is not necessary for such fields to have the same symmetries as the background spacetime. On the other hand, if one considers the backreaction of the matter fields over the spacetime, in order to look for exact solutions to the field equations, the relation between symmetries of the spacetime and the matter fields is not trivial, for a recent study about symmetry inheritance of scalar fields see [54] and references therein. In the case considered here, the gravitational field equations imply that the trace of the stress-energy tensor must vanish, due to the Bach tensor is traceless, therefore if one go beyond the probe-field approximation, this implies that the stress-energy tensor of the matter fields must be traceless. Based on these arguments, first we will consider a test scalar field minimally coupled to curvature, then we will consider a test scalar field conformally coupled to curvature, which have a traceless stress-energy tensor, and we find analytically and numerically the quasinormal frequencies for scalar fields for some special cases.

A. Scalar field minimally coupled to gravity

In this section we calculate the QNMs of the $z = 0$ Lifshitz black hole for a test scalar field minimally coupled to gravity. The Klein-Gordon equation in curved spacetime is

$$\frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g} g^{\mu\nu} \partial_\nu) \psi = m^2 \psi , \quad (9)$$

where m is the mass of the scalar field ψ , which is minimally coupled to curvature. By means of the following ansatz

$$\psi = e^{-i\omega t} R(r) Y(\theta, \phi) , \quad (10)$$

where $Y(\theta, \phi)$ is a normalizable harmonic function on the two-sphere which satisfies

$$\nabla^2 Y = -\kappa Y , \quad (11)$$

being $\kappa = l(l+1)$ the eigenvalues for the spheric manifold, with $l = 0, 1, 2, \dots$, the Klein-Gordon equation reduces to

$$\frac{1}{4r} \partial_r (r^3 f(r) \partial_r R) + \left(\frac{\omega^2 \ell^2}{f(r)} - \frac{\kappa \ell^2}{r^2} - m^2 \ell^2 \right) R(r) = 0 . \quad (12)$$

Now, by considering $R(r) = K(r)/r$ and by introducing the tortoise coordinate r_* , given by $dr_* = \frac{2\ell dr}{rf(r)}$, the latter equation can be rewritten as a one-dimensional Schrödinger equation

$$[\partial_{r_*}^2 + \omega^2 - V_{eff}(r)] K(r_*) = 0 , \quad (13)$$

where the effective potential is given by

$$V_{eff}(r) = \frac{f(r)}{4} \left[\frac{f(r)}{\ell^2} + \frac{rf'(r)}{\ell^2} + \frac{4\kappa}{r^2} + 4m^2 \right] . \quad (14)$$

In Fig. (1) we plot the effective potential for $\kappa = 2$ and in Fig. (2) for $\kappa = 0$ and different values of the parameter λ . Note that when $r \rightarrow \infty$ the effective potential goes to $1/(4\ell^2) + m^2$.




FIG. 1: The effective potential as a function of r , for $m = 1$, $\ell = 1$ and $\kappa = 2$.

1. Case $\kappa = 0$

In order to find analytical solutions to the radial equation (12), we perform the change of variables $y = r^2$ and get the following equation:

$$y(y - y_+)(y - y_-) R''(y) + y(2y + \lambda) R'(y) + \left(\frac{\omega^2 \ell^2 y^3}{(y - y_+)(y - y_-)} - \kappa \ell^2 - m^2 \ell^2 y \right) R(y) = 0 , \quad (15)$$




FIG. 2: The effective potential as a function of r , for $m = 1$, $\ell = 1$ and $\kappa = 0$.

where the prime denotes the derivative with respect to y , and y_+ and y_- are the roots of

$$f(y) = 1 + \frac{\lambda}{y} + \frac{\lambda^2 - \ell^4}{3y^2}, \quad (16)$$

and are given by

$$y_{\pm} = -\frac{\lambda}{2} \pm \sqrt{-\frac{\lambda^2}{12} + \frac{\ell^4}{3}}. \quad (17)$$

Additionally, performing another change of variable $z = 1 - \frac{y_+}{y}$ and noting that $\lambda = -(y_+ + y_-)$, we arrive at the following expression

$$z(z-1)(z-(1-Q))R''(z) - (z-1)(1-Q-2z)R'(z) + \left(\frac{\omega^2 \ell^2 Q^2}{z(z-1)(z-(1-Q))} + \frac{\kappa \ell^2}{y_-} - \frac{m^2 \ell^2 Q}{z-1} \right) R(z) = 0, \quad (18)$$

where we have defined $Q = y_+/y_-$ and now a prime means derivative with respect to z . In Fig. (3), we plot Q as a function of λ/ℓ^2 and we observe that Q can be positive or negative depending on the values of the parameter λ :

$$Q > 1 \text{ for } -2\ell^2 \leq \lambda < -\ell^2, \quad (19)$$

$$Q < 0 \text{ for } -\ell^2 < \lambda < \ell^2. \quad (20)$$

On the other hand, equation (18) can be manipulated and put into the following form


$$R''(z) + \left(\frac{1}{z} + \frac{1}{z-(1-Q)} \right) R'(z) + \left(\frac{\omega^2 \ell^2 Q^2 / (1-Q)}{z} + \frac{Q(\omega^2 \ell^2 - m^2 \ell^2)}{z-1} + \frac{\kappa \ell^2}{y_-} - \frac{Q\omega^2 \ell^2 / (1-Q)}{z-(1-Q)} \right) \frac{1}{z(z-1)(z-(1-Q))} R(z) = 0. \quad (21)$$

We note that for $\kappa = 0$ this equation corresponds to a Riemann differential equation, whose general form is [55]

$$\frac{d^2 w}{dz^2} + \left(\frac{1-\alpha-\alpha'}{z-a} + \frac{1-\beta-\beta'}{z-b} + \frac{1-\gamma-\gamma'}{z-c} \right) \frac{dw}{dz} + \left(\frac{\alpha\alpha'(a-b)(a-c)}{z-a} + \frac{\beta\beta'(b-c)(b-a)}{z-b} + \frac{\gamma\gamma'(c-a)(c-b)}{z-c} \right) \frac{w}{(z-a)(z-b)(z-c)} = 0, \quad (22)$$

where a, b and c are the singular points, and the exponents $\alpha, \alpha', \beta, \beta', \gamma$ and γ' are subject to the condition

$$\alpha + \alpha' + \beta + \beta' + \gamma + \gamma' = 1. \quad (23)$$

FIG. 3: Q as a function of λ/ℓ^2 .

The complete solution is denoted by the symbol

$$w = P \left\{ \begin{matrix} a & b & c \\ \alpha & \beta & \gamma \\ \alpha' & \beta' & \gamma' \end{matrix} \middle| z \right\} , \quad (24)$$

and the Riemann P function can be reduced to the hypergeometric function through

$$w = \left(\frac{z-a}{z-b} \right)^\alpha \left(\frac{z-c}{z-b} \right)^\gamma P \left\{ \begin{matrix} 0 & \infty & 1 \\ 0 & \alpha + \beta + \gamma & 0 \\ \alpha' - \alpha & \alpha + \beta' + \gamma & \gamma' - \gamma \end{matrix} \middle| \frac{(z-a)(c-b)}{(z-b)(c-a)} \right\} , \quad (25)$$

where the P function is now Gauss hypergeometric function [55]. We observe that, in the radial equation (21), the regular singular points a, b and c have the values

$$a = 0 , \quad b = 1 - Q , \quad c = 1 , \quad (26)$$

and the exponents are given by

$$\alpha = \pm \frac{i\omega\ell y_+}{y_+ - y_-} , \alpha' = \mp \frac{i\omega\ell y_+}{y_+ - y_-} , \quad (27)$$

$$\beta = \pm \frac{i\omega\ell |y_-|}{y_+ - y_-} , \beta' = \mp \frac{i\omega\ell |y_-|}{y_+ - y_-} , \quad (28)$$

$$\gamma = \frac{1}{2} \pm \sqrt{\frac{1}{4} - (\omega^2 - m^2)\ell^2} , \gamma' = \frac{1}{2} \mp \sqrt{\frac{1}{4} - (\omega^2 - m^2)\ell^2} . \quad (29)$$

Therefore, the solution to equation (21) can be written as

$$R(z) = C_1 \left(\frac{z}{z-1+Q} \right)^\alpha \left(\frac{z-1}{z-1+Q} \right)^\gamma {}_2F_1 \left(A, B, C, \frac{Qz}{z-(1-Q)} \right) + \\ C_2 \left(\frac{z}{z-1+Q} \right)^{\alpha'} \left(\frac{z-1}{z-1+Q} \right)^\gamma {}_2F_1 \left(A-C+1, B-C+1, 2-C, \frac{Qz}{z-(1-Q)} \right) , \quad (30)$$

where we have defined the constants A, B and C as

$$\begin{aligned} A &= \alpha + \beta + \gamma , \\ B &= \alpha + \beta' + \gamma , \\ C &= 1 + \alpha - \alpha' . \end{aligned} \quad (31)$$

In the near-horizon limit, the above expression behaves as

$$R(z \rightarrow 0) = \frac{(-1)^\gamma C_1}{(-1+Q)^{\alpha+\gamma}} z^\alpha + \frac{(-1)^\gamma C_2}{(-1+Q)^{\alpha'+\gamma}} z^{\alpha'} , \quad (32)$$

Now, we impose as a boundary condition that classically nothing can escape from the event horizon. So, choosing the exponent α as

$$\alpha = -\frac{i\omega\ell y_+}{y_+ - y_-} , \quad (33)$$

implies that we must take $C_2 = 0$ in order to have only ingoing waves at the horizon. Therefore, our solution simplifies to

$$R(z) = C_1 \left(\frac{z}{z-1+Q} \right)^\alpha \left(\frac{z-1}{z-1+Q} \right)^\gamma {}_2F_1 \left(A, B, C, \frac{Qz}{z-(1-Q)} \right) . \quad (34)$$

Now, we implement boundary conditions at spatial infinity. In order to do so we employ the Kummer relations [55], and write the solution as

$$\begin{aligned} R(z) = & C_1 \left(\frac{z}{z-1+Q} \right)^\alpha \left(\frac{z-1}{z-1+Q} \right)^\gamma \frac{\Gamma(C)\Gamma(C-A-B)}{\Gamma(C-A)\Gamma(C-B)} {}_2F_1 \left(A, B, A+B-C, 1 - \frac{Qz}{z-(1-Q)} \right) + \\ & C_1 (1-Q)^{\gamma'-\gamma} \left(\frac{z}{z-1+Q} \right)^\alpha \left(\frac{z-1}{z-1+Q} \right)^{\gamma'} \frac{\Gamma(C)\Gamma(A+B-C)}{\Gamma(A)\Gamma(B)} \times \\ & {}_2F_1 \left(C-A, C-B, C-A-B+1, 1 - \frac{Qz}{z-(1-Q)} \right) . \end{aligned} \quad (35)$$

At the limit $z \rightarrow 1$, the above solution becomes

$$R(z \rightarrow 1) = \frac{C_1}{Q^{\alpha+\gamma}} (z-1)^\gamma \frac{\Gamma(C)\Gamma(C-A-B)}{\Gamma(C-A)\Gamma(C-B)} + \frac{C_1 (1-Q)^{\gamma'-\gamma}}{Q^{\alpha+\gamma'}} (z-1)^{\gamma'} \frac{\Gamma(C)\Gamma(A+B-C)}{\Gamma(A)\Gamma(B)} . \quad (36)$$

Now, we choose the exponents γ and γ' as follows

$$\begin{aligned} \gamma &= \frac{1}{2} + \sqrt{\frac{1}{4} - (\omega^2 - m^2)\ell^2} , \\ \gamma' &= \frac{1}{2} - \sqrt{\frac{1}{4} - (\omega^2 - m^2)\ell^2} . \end{aligned} \quad (37)$$

So, imposing the condition that the scalar field be null at spatial infinity, we can determine the QNFs. The second term of equation (36) blows up when $z \rightarrow 1$ unless we impose the condition $A = -n$ or $B = -n$; therefore, we obtain the following set of QNFs:

$$\omega\ell = \frac{i(m^2\ell^2 - n(1+n))}{1+2n} . \quad (38)$$

These QNFs are purely imaginary and negative for $m = 0$, which guarantees that the Lifshitz black hole is stable under massless scalar field perturbations for the mode with the lowest angular momentum. For $m > 0$ there are QNFs with imaginary and positive value, and the Lifshitz black hole is unstable under scalar field perturbations. Also, we note that if we interchange the values of the exponents in equation (37) the same QNFs are obtained. It is worth mentioning that Eq. (15) with $\kappa = 0$ can be written as

$$z(1-z)R''(z) + (1-z)R'(z) + \left(\frac{\omega^2\ell^2(z y_- - y_+)^2}{(y_+ - y_-)^2 z(1-z)} - \frac{m^2\ell^2}{1-z} \right) R(z) = 0 , \quad (39)$$

under the change of variable $z = \frac{y_- y_+}{y - y_-}$, and if we define $R(z) = z^\alpha (1-z)^\beta F(z)$, the above equation leads to the hypergeometric equation

$$z(1-z)F''(z) + [c - (1+a+b)z]F'(z) - abF(z) = 0 , \quad (40)$$

where

$$\alpha = \pm \frac{i\omega\ell y_+}{y_+ - y_-} , \quad (41)$$

$$\beta = \frac{1}{2} \left(1 \pm \sqrt{1 + 4(m^2 - \omega^2)\ell^2} \right) , \quad (42)$$

and the constants are given by

$$a = \alpha + \beta - \frac{i\omega\ell |y_-|}{(y_+ - y_-)} , \quad (43)$$

$$b = \alpha + \beta + \frac{i\omega\ell |y_-|}{(y_+ - y_-)} , \quad (44)$$

$$c = 1 + 2\alpha . \quad (45)$$

The general solution of the hypergeometric equation (50) is

$$F(z) = c_1 {}_2F_1(a, b, c; z) + c_2 z^{1-c} {}_2F_1(a - c + 1, b - c + 1, 2 - c; z) , \quad (46)$$

and it has three regular singular points at $z = 0$, $z = 1$, and $z = \infty$. ${}_2F_1(a, b, c; z)$ is a hypergeometric function and c_1 and c_2 are integration constants. Note that the above QNFs could be computed using the solution (46).

2. Case $Q = \pm\infty$

In this case it is possible to obtain an analytical solution for all values of the angular momentum κ . Thus, for $\lambda = -\ell^2$ or equivalently $Q = \pm\infty$, the radial equation (12) can be written as

$$z(1-z)\partial_z^2 R(z) + (1-z)\partial_z R(z) + \left[\frac{\omega^2\ell^2}{z(1-z)} - \frac{m^2\ell^2}{1-z} - \kappa \right] R(z) = 0 , \quad (47)$$

where we have considered $z = 1 - \ell^2/r^2$. Using the decomposition $R(z) = z^\alpha(1-z)^\beta K(z)$, with

$$\alpha_\pm = \pm i\omega\ell , \quad (48)$$

$$\beta_\pm = \frac{1}{2} \left(1 \pm \sqrt{1 + 4(m^2 - \omega^2)\ell^2} \right) , \quad (49)$$

we can write (47) as a hypergeometric equation for K

$$z(1-z)K''(z) + [c - (1+a+b)z]K'(z) - abK(z) = 0 , \quad (50)$$

where the coefficients are given by

$$a = \alpha + \beta \mp \sqrt{-\kappa} , \quad (51)$$

$$b = \alpha + \beta \pm \sqrt{-\kappa} , \quad (52)$$

$$c = 1 + 2\alpha . \quad (53)$$

The general solution of the hypergeometric equation (50) is

$$K = C_1 {}_2F_1(a, b, c; z) + C_2 z^{1-c} {}_2F_1(a - c + 1, b - c + 1, 2 - c; z) , \quad (54)$$

and it has three regular singular points at $z = 0$, $z = 1$, and $z = \infty$. ${}_2F_1(a, b, c; z)$ is a hypergeometric function and C_1 and C_2 are constants. Thus, the solution for the radial function $R(z)$ is

$$R(z) = C_1 z^\alpha (1-z)^\beta {}_2F_1(a, b, c; z) + C_2 z^{-\alpha} (1-z)^\beta {}_2F_1(a-c+1, b-c+1, 2-c; z) . \quad (55)$$

So, in the vicinity of the horizon, $z = 0$ and using the property $F(a, b, c, 0) = 1$, the function $R(z)$ behaves as

$$R(z) = C_1 e^{\alpha \ln z} + C_2 e^{-\alpha \ln z} , \quad (56)$$

and the scalar field ψ , for $\alpha = \alpha_-$, can be written as follows:

$$\psi \sim C_1 e^{-i\omega\ell(t+\ln z)} + C_2 e^{-i\omega\ell(t-\ln z)} , \quad (57)$$

in which, the first term represents an ingoing wave and the second an outgoing wave in the black hole. So, by imposing that only ingoing waves existing at the horizon, this fixes $C_2 = 0$. The radial solution then becomes

$$R(z) = C_1 e^{\alpha \ln z} (1-z)^\beta {}_2F_1(a, b, c; z) = C_1 e^{-i\omega\ell \ln z} (1-z)^\beta {}_2F_1(a, b, c; z) . \quad (58)$$

To implement boundary conditions at infinity ($z = 1$), we apply Kummer's formula for the hypergeometric function [55],

$${}_2F_1(a, b, c; z) = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)} F_1 + (1-z)^{c-a-b} \frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)} F_2 , \quad (59)$$

where,

$$F_1 = {}_2F_1(a, b, a+b-c, 1-z) , \quad (60)$$

$$F_2 = {}_2F_1(c-a, c-b, c-a-b+1, 1-z) . \quad (61)$$

With this expression, the radial function (58) reads

$$R(z) = C_1 e^{-i\omega\ell \ln z} (1-z)^\beta \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)} F_1 + C_1 e^{-i\omega\ell \ln z} (1-z)^{c-a-b+\beta} \frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)} F_2 , \quad (62)$$

and at infinity it can be written as

$$R_{asympt.}(z) = C_1 (1-z)^\beta \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)} + C_1 (1-z)^{1-\beta} \frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)} . \quad (63)$$

So, for $\beta_+ > 1$, the field at infinity vanishes if $(a)|_{\beta_+} = -n$ or $(b)|_{\beta_+} = -n$ for $n = 0, 1, 2, \dots$, and for $\beta_- < 0$, the field at infinity vanishes if $(c-a)|_{\beta_-} = -n$ or $(c-b)|_{\beta_-} = -n$. Therefore, the QNFs are given by

$$\omega\ell = -i \frac{-m^2\ell^2 + n + n^2 + \kappa \mp \sqrt{-\kappa}(1+2n)}{1+2n \mp 2\sqrt{-\kappa}} , \quad (64)$$

where $\sqrt{-\kappa} = i\sqrt{l(l+1)}$. This expression can be written as ¹

$$\omega\ell = \pm \frac{\sqrt{\kappa}(-(1+2n)^2 + 2(-m^2\ell^2 + n + n^2 - \kappa))}{(1+2n)^2 + 4\kappa} - i \frac{(1+2n)(-m^2\ell^2 + n + n^2 + \kappa)}{(1+2n)^2 + 4\kappa} . \quad (65)$$

Because not all the QNFs have a negative imaginary part we conclude that this black hole is not stable under scalar field perturbations for the case when $Q = \pm\infty$.

¹ The same QNFs can be obtained by imposing that only outgoing waves exist at spatial infinity.

3. Case $Q = 1$

In the extremal case $\lambda = -2\ell^2$, or equivalently $Q = 1$, the radial equation (18) reads

$$z(z-1)R''(z) + 2(z-1)R'(z) + \left(\frac{\omega^2\ell^2}{z^3(z-1)} + \frac{\kappa}{z} - \frac{m^2\ell^2}{z(z-1)} \right) R(z) = 0, \quad (66)$$

and its solution is given by

$$\begin{aligned} R(z) = & C_1 e^{\frac{i\omega\ell}{z}} Heun_C \left(-2i\omega\ell, \sqrt{1-4(\omega^2-m^2)\ell^2}, 1, -2\omega^2\ell^2, -\kappa + \frac{1}{2} + 2\omega^2\ell^2, \frac{z-1}{z} \right) z^{-\frac{3}{2}-\frac{1}{2}\sqrt{1-4(\omega^2-m^2)\ell^2}} \times \\ & (z-1)^{\frac{1}{2}+\frac{1}{2}\sqrt{1-4(\omega^2-m^2)\ell^2}} \\ & + C_2 e^{\frac{i\omega\ell}{z}} Heun_C \left(-2i\omega\ell, -\sqrt{1-4(\omega^2-m^2)\ell^2}, 1, -2\omega^2\ell^2, -\kappa + \frac{1}{2} + 2\omega^2\ell^2, \frac{z-1}{z} \right) \times \\ & z^{-\frac{3}{2}+\frac{1}{2}\sqrt{1-4(\omega^2-m^2)\ell^2}} (z-1)^{\frac{1}{2}-\frac{1}{2}\sqrt{1-4(\omega^2-m^2)\ell^2}}, \end{aligned} \quad (67)$$

where $Heun_C$ is the confluent Heun function. Thus, when $z \rightarrow 1$, and in order to have a regular scalar field at spatial infinity, we must set $C_2 = 0$; therefore, the solution reduces to

$$\begin{aligned} R(z) = & C_1 e^{\frac{i\omega\ell}{z}} Heun_C \left(-2i\omega\ell, \sqrt{1-4(\omega^2-m^2)\ell^2}, 1, -2\omega^2\ell^2, -\kappa + \frac{1}{2} + 2\omega^2\ell^2, \frac{z-1}{z} \right) z^{-\frac{3}{2}-\frac{1}{2}\sqrt{1-4(\omega^2-m^2)\ell^2}} \times \\ & (z-1)^{\frac{1}{2}+\frac{1}{2}\sqrt{1-4(\omega^2-m^2)\ell^2}}, \end{aligned} \quad (68)$$

where the property $Heun_C(a, b, c, d, e, 0) = 1$ was used [56]. However, we observe that when $z \rightarrow 1$, the scalar field is null $R(z) \rightarrow 0$; therefore, there are no QNMs in this case.

B. Scalar field conformally coupled to gravity

In this section we calculate the QNMs of the $z = 0$ Lifshitz black hole for a test scalar field conformally coupled to gravity. The Klein-Gordon equation for a scalar field non-minimally coupled to curvature is

$$\frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g} g^{\mu\nu} \partial_\nu) \psi - \xi \mathcal{R} \psi = m^2 \psi, \quad (69)$$

where m is the mass of the scalar field ψ , ξ is the non-minimal coupling parameter and \mathcal{R} is the Ricci scalar, which reads

$$\mathcal{R} = \frac{\ell^4 - \lambda^2}{2\ell^2 r^4} + \frac{4\ell^2 - \lambda}{2\ell^2 r^2} - \frac{3}{2\ell^2}. \quad (70)$$

For a conformally coupled scalar field case we must take $m = 0$ and $\xi = 1/6$. Now, by means of the following ansatz

$$\psi = e^{-i\omega t} R(r) Y(\theta, \phi), \quad (71)$$

where $Y(\theta, \phi)$ is a normalizable harmonic function on the two-sphere which satisfies Eq. (11), the Klein-Gordon equation reduces to

$$\frac{1}{4r} \partial_r (r^3 f(r) \partial_r R) + \left(\frac{\omega^2\ell^2}{f(r)} - \frac{\kappa\ell^2}{r^2} - m^2\ell^2 - \xi\ell^2\mathcal{R} \right) R(r) = 0, \quad (72)$$

which can be written as a one-dimensional Schrödinger equation with an effective potential that vanishes at spatial infinity. Therefore, we will consider only outgoing waves at the asymptotic region as boundary condition. It is worth to mention that Eq. (72) only has analytical solution for $\lambda = -1$ as we will show below. Therefore, we will perform numerical studies for $\lambda \neq -1$ by using the improved AIM [57], which is an improved version of the method proposed in Refs. [58, 59] and it has been applied successful in the context of QNMs for different black holes geometries, see for instance [41, 57, 60–62].

1. Numerical analysis

In order to implement the improved AIM we make the following consecutive change of variables $y = r^2$ and $z = \frac{y-y_+}{y-y_-}$ to Eq. (69), as we do in the previous sections. Then, the Klein-Gordon equation yields

$$z(1-z)R''(z) + (1-z)R'(z) + \left(\frac{\omega^2 \ell^2 (zy_- - y_+)^2}{(y_+ - y_-)^2 z(1-z)} + \frac{\kappa \ell^2}{zy_- - y_+} - \frac{\xi(\ell^4 - \lambda^2)(1-z)}{2(zy_- - y_+)^2} + \frac{\xi(4\ell^2 - \lambda)}{2(zy_- - y_+)} + \frac{3\xi}{2(1-z)} \right) R(z) = 0. \quad (73)$$

Now, we must consider the behavior of the scalar field on the event horizon and at spatial infinity. Accordingly, on the horizon, $z \rightarrow 0$, the behavior of the scalar field is given by

$$R(z \rightarrow 0) \sim C_1 z^{-\frac{i\omega \ell y_+}{y_+ - y_-}} + C_2 y^{\frac{i\omega \ell y_+}{y_+ - y_-}}, \quad (74)$$

So, if we consider only ingoing waves on the horizon, we must impose $C_2 = 0$. Asymptotically, from Eq. (73), the scalar field behaves as

$$R(z \rightarrow 1) \sim D_1 (1-z)^{1/2-i\omega \ell} + D_2 (1-z)^{1/2+i\omega \ell}. \quad (75)$$

So, in order to have only outgoing waves at infinity we must impose $D_2 = 0$. Therefore, taking into account these behaviors we define

$$R(z) = z^{-\frac{i\omega \ell y_+}{y_+ - y_-}} (1-z)^{1/2-i\omega \ell} \chi(z). \quad (76)$$

Then, by inserting these fields in Eq. (73) we obtain the homogeneous linear second-order differential equation for the function $\chi(z)$

$$\chi'' = \lambda_0(z)\chi' + s_0(z)\chi, \quad (77)$$

where

$$\lambda_0(z) = -\frac{(y_+ - y_-)(1-2z) - 2i\omega \ell(y_+ + y_-z - 2y_+z)}{(y_+ - y_-)z(1-z)}, \quad (78)$$

$$s_0(z) = \frac{\ell^4 + 3y_+^2 + z(-\ell^4 + y_-(-4\ell^2 + 3y_-z - 12\kappa\ell^2)) + \lambda y_-z - \lambda^2(1-z) - y_-(-4\ell^2 + 6y_-z - 12\kappa\ell^2 + \lambda)}{12z(1-z)(y_+ - y_-z)^2} \quad (79)$$

$$+ \frac{12i\omega \ell(y_- - 2y_+)(y_+ - y_-z)^2 - 48\omega^2 \ell^2 y_+(y_+ - y_-z)^2}{12z(1-z)(y_+ - y_-)(y_+ - y_-z)^2}.$$

Then, in order to implement the improved AIM it is necessary to differentiate Eq. (77) n times with respect to z , which yields the following equation:

$$\chi^{n+2} = \lambda_n(z)\chi' + s_n(z)\chi, \quad (80)$$

where

$$\lambda_n(z) = \lambda'_{n-1}(z) + s_{n-1}(z) + \lambda_0(z)\lambda_{n-1}(z), \quad (81)$$

$$s_n(z) = s'_{n-1}(z) + s_0(z)\lambda_{n-1}(z). \quad (82)$$

Then, by expanding the λ_n and s_n in a Taylor series around the point δ , at which the improved AIM is performed

$$\lambda_n(\delta) = \sum_{i=0}^{\infty} c_n^i (z - \delta)^i, \quad (83)$$

$$s_n(\delta) = \sum_{i=0}^{\infty} d_n^i (z - \delta)^i, \quad (84)$$

where the c_n^i and d_n^i are the i^{th} Taylor coefficients of $\lambda_n(\delta)$ and $s_n(\delta)$, respectively, and by replacing the above expansions in Eqs. (81) and (82) the following set of recursion relations for the coefficients is obtained:

$$c_n^i = (i+1)c_{n-1}^{i+1} + d_{n-1}^i + \sum_{k=0}^i c_0^k c_{n-1}^{i-k} , \quad (85)$$

$$d_n^i = (i+1)d_{n-1}^{i+1} + \sum_{k=0}^i d_0^k c_{n-1}^{i-k} . \quad (86)$$

In this manner, the authors of the improved AIM have avoided the derivatives that contain the AIM in [57, 60], and the quantization condition, which is equivalent to imposing a termination to the number of iterations, is given by

$$d_n^0 c_{n-1}^0 - d_{n-1}^0 c_n^0 = 0 . \quad (87)$$

We solve this equation numerically to find the QNFs. In Table I, we show some lowest QNFs, for a scalar field conformally coupled to curvature with $\kappa = 0$, $\ell = 1$ and different values of λ . Then, in Table II, we show some fundamentals QNFs, for a scalar field conformally coupled to curvature with $\kappa = 0, 1, 2, 3$, $\ell = 1$ and different values of λ . We observe that the QNFs have real and imaginary parts, with an imaginary part that is negative, which ensures the stability of the 4-dimensional Lifshitz black hole under scalar perturbations.

TABLE I: Quasinormal frequencies for $\kappa = 0$, $\ell = 1$ and different values of λ .

n	$\lambda = -1.3$	$\lambda = -1$	$\lambda = -0.7$
0	0.29170 - 0.22801 <i>i</i>	0.32275 - 0.25000 <i>i</i>	0.35757 - 0.27902 <i>i</i>
1	0.27600 - 0.68476 <i>i</i>	0.32275 - 0.75000 <i>i</i>	0.34211 - 0.83763 <i>i</i>
2	0.23817 - 1.14518 <i>i</i>	0.32275 - 1.25000 <i>i</i>	0.30647 - 1.39878 <i>i</i>
3	0.13120 - 1.63955 <i>i</i>	0.32275 - 1.75000 <i>i</i>	0.22919 - 1.97158 <i>i</i>

TABLE II: Fundamentals Quasinormal frequencies ($n = 0$), $\ell = 1$ and different values of κ and λ .

κ	$\lambda = -1.9$	$\lambda = -1.3$	$\lambda = -1$	$\lambda = -0.7$	$\lambda = -0.3$
0	0.22840 - 0.20893 <i>i</i>	0.29170 - 0.22801 <i>i</i>	0.32275 - 0.25000 <i>i</i>	0.35757 - 0.27902 <i>i</i>	0.41878 - 0.33808 <i>i</i>
2	0.60092 - 0.19838 <i>i</i>	0.70778 - 0.22800 <i>i</i>	0.77728 - 0.25000 <i>i</i>	0.86642 - 0.27902 <i>i</i>	1.04077 - 0.33703 <i>i</i>
6	0.98966 - 0.19724 <i>i</i>	1.15443 - 0.22800 <i>i</i>	1.26656 - 0.25000 <i>i</i>	1.41292 - 0.27902 <i>i</i>	1.70231 - 0.33686 <i>i</i>
12	1.38091 - 0.19692 <i>i</i>	1.60635 - 0.22800 <i>i</i>	1.76186 - 0.25000 <i>i</i>	1.96592 - 0.27902 <i>i</i>	2.37061 - 0.33682 <i>i</i>

2. Case $Q = \pm\infty$

In this case ($\lambda = -\ell^2$), due to the simplicity of the Ricci scalar, it is possible to obtain an analytical solution. The radial equation (72) reads

$$\frac{1}{4r} \partial_r (r^3 f(r) \partial_r R) + \left(\frac{\omega^2 \ell^2}{f(r)} - \frac{\kappa \ell^2 + 5\xi \ell^2/2}{r^2} - m^2 \ell^2 + \frac{3\xi}{2} \right) R(r) = 0 . \quad (88)$$

So, if we compare this equation with the analogous equation of the minimal case (Eq. (12)), we see that is possible to obtain (88) by means of the following substitutions in Eq. (12):

$$\begin{aligned} \kappa &\rightarrow \kappa + \frac{5\xi}{2} , \\ m^2 &\rightarrow m^2 - \frac{3\xi}{2\ell^2} . \end{aligned} \quad (89)$$

Thus, using the above substitutions in the QNFs (65) we find

$$\omega \ell = \pm \frac{\sqrt{\kappa + \frac{5\xi}{2}} \left(-(1+2n)^2 + 2(-m^2 \ell^2 + n + n^2 - \kappa - \xi) \right)}{(1+2n)^2 + 4\kappa + 10\xi} - i \frac{(1+2n)(-m^2 \ell^2 + n + n^2 + \kappa + 4\xi)}{(1+2n)^2 + 4\kappa + 10\xi} , \quad (90)$$

and for a conformal scalar field ($m = 0$, $\xi = 1/6$) this equation yields

$$\omega\ell = \pm \frac{\sqrt{\kappa + \frac{5}{12}} \left(-(1+2n)^2 + 2(n+n^2 - \kappa - \frac{1}{6}) \right)}{(1+2n)^2 + 4\kappa + \frac{5}{3}} - i \frac{(1+2n)(n+n^2 + \kappa + \frac{2}{3})}{(1+2n)^2 + 4\kappa + \frac{5}{3}}, \quad (91)$$

which satisfies that there are only outgoing waves at the asymptotic region, see footnote 1. Clearly the imaginary part of the QNFs is negative, which ensures the stability of this black hole under conformally coupled scalar field perturbations. These QNFs agrees with the numerical results for $\lambda = -1$, $\kappa = 0$ and $\ell = 1$ showed in Table (I).

IV. REFLECTION AND TRANSMISSION COEFFICIENTS AND ABSORPTION CROSS SECTION

In this section, we focus our attention to the minimally coupled scalar field case. However, a similar analysis can be performed for scalar fields conformally coupled to gravity, and for $\lambda = -\ell^2$ the results are straightforward obtained from the minimal case by using the substitutions (89) and taking $m = 0$, $\xi = 1/6$. The reflection and the transmission coefficients are defined by

$$\Re := \left| \frac{F_{\text{asympt}}^{\text{out}}}{F_{\text{asympt}}^{\text{in}}} \right|, \quad \text{and} \quad \Im := \left| \frac{F_{\text{hor}}^{\text{in}}}{F_{\text{asympt}}^{\text{in}}} \right|, \quad (92)$$

where F is the flux given by

$$F = \frac{1}{2i} \sqrt{-g} g^{rr} (R^* \partial_r R - R \partial_r R^*). \quad (93)$$

So, in order to calculate the above coefficients we need to know the behavior of the radial function both on the horizon and at asymptotic infinity.

A. Case $\kappa = 0$

In this case, the behavior at the horizon is given by Eq. (46) with $c_2 = 0$, and choosing the negative value of α and using Eq. (93), we get the following flux on the horizon:

$$F_{\text{hor}}^{\text{in}} = -\omega \sin \theta |c_1|^2 y_+. \quad (94)$$

On the other hand, by applying Kummer's formula (59) for the hypergeometric function in Eq. (46), the asymptotic behavior of $R(z)$ can be written as

$$R(z \rightarrow 1) = \bar{c}_1 (z-1)^\beta + \bar{c}_2 (z-1)^{1-\beta}, \quad (95)$$

where

$$\begin{aligned} \bar{c}_1 &= c_1 \frac{\Gamma(c) \Gamma(c-a-b)}{\Gamma(c-a) \Gamma(c-b)}, \\ \bar{c}_2 &= c_1 \frac{\Gamma(c) \Gamma(a+b-c)}{\Gamma(a) \Gamma(b)}. \end{aligned} \quad (96)$$

Thus, using Eq. (93) we obtain the flux

$$F_{\text{asympt.}} = \frac{\sin \theta}{\ell} (y_+ - y_-) \sqrt{(\omega^2 - m^2) \ell^2 - \frac{1}{4}} \left(|\bar{c}_2|^2 - |\bar{c}_1|^2 \right), \quad (97)$$

for $\beta = \beta_+$, the reflection and transmission coefficients are given by

$$\Re = \frac{|\bar{c}_2|^2}{|\bar{c}_1|^2}, \quad (98)$$

$$\Im = \frac{\omega\ell |c_1|^2 y_+}{(y_+ - y_-) |\bar{c}_1|^2 \sqrt{(\omega^2 - m^2) \ell^2 - \frac{1}{4}}}, \quad (99)$$




FIG. 4: The reflection coefficient R (solid curve), the transmission coefficient T (dashed curve), $R + T$ (thick curve) and the absorption cross section σ_{abs} (dotted curve) as a function of ω , for $m = 1$, $\ell = 1$, and $\lambda = -1.9$.




FIG. 5: The behavior of σ_{abs} as a function of ω , for $\lambda = -1.9$, $\ell = 1$, and $m = 1, 1.5, 2$.

and the absorption cross section, σ_{abs} , is given by

$$\sigma_{abs} = \frac{\ell |c_1|^2 y_+}{(y_+ - y_-) |\bar{c}_1|^2 \sqrt{(\omega^2 - m^2) \ell^2 - \frac{1}{4}}} . \quad (100)$$

Interestingly, the poles of the transmission coefficient yields the same set of QNFs found in the previous section, which is equivalent to imposing as a boundary condition that only outgoing waves exist at asymptotic infinity. Now, we perform a numerical analysis of the reflection coefficient (98), transmission coefficient (99) and absorption cross section (100) of the four-dimensional Lifshitz black hole with $z = 0$ for scalar fields. So, we plot the reflection and transmission coefficients and the absorption cross section in Fig. (4) for scalar fields with $m = 1$. Essentially, we found that the reflection coefficient is 1 at low frequency limit, and for high frequency limit this coefficient is null, with the behavior of the transmission coefficient being opposite with $R + T = 1$. In addition, the absorption cross section is null in the low and high-frequency limit, but there is a range of frequencies for which the absorption cross section is not null, and it also has a maximum value in the low-frequency limit (see Fig. (5)). Furthermore, we observe that the absorption cross section can take higher values when the mass of the scalar field decreases (Fig. (5)) in the low frequency limit. However, beyond a certain value of the frequency the absorption cross section does not depend on the mass of the scalar field.

B. Case $Q = \pm\infty$

In this case the behavior at the horizon is given by Eq. (56), with $C_2 = 0$, and using Eq. (93), we get the flux at the horizon

$$F_{hor}^{in} = -\omega \ell^2 \sin \theta |C_1|^2 . \quad (101)$$

On the other hand, the asymptotic behavior of $R(z)$ is given by Eq. (63), which can be written as

$$R(z \rightarrow 1) = B_1(z-1)^\beta + B_2(z-1)^{1-\beta}, \quad (102)$$

where

$$\begin{aligned} B_1 &= C_1 \frac{\Gamma(c) \Gamma(c-a-b)}{\Gamma(c-a) \Gamma(c-b)}, \\ B_2 &= C_1 \frac{\Gamma(c) \Gamma(a+b-c)}{\Gamma(a) \Gamma(b)}. \end{aligned} \quad (103)$$

Thus, using Eq. (93), we get the flux

$$F_{asymp.} = \ell \sin \theta \sqrt{(\omega^2 - m^2) \ell^2 - \frac{1}{4} (|B_2|^2 - |B_1|^2)} \quad (104)$$

Therefore, the reflection and transmission coefficients are given by

$$\mathfrak{R} = \frac{|B_2|^2}{|B_1|^2}, \quad (105)$$

$$\mathfrak{T} = \frac{\omega \ell |C_1|^2}{\sqrt{(\omega^2 - m^2) \ell^2 - \frac{1}{4} |B_1|^2}}, \quad (106)$$

and the absorption cross section, σ_{abs} , is given by

$$\sigma_{abs} = \frac{\mathfrak{T}}{\omega} = \frac{\ell |C_1|^2}{\sqrt{(\omega^2 - m^2) \ell^2 - \frac{1}{4} |B_1|^2}}. \quad (107)$$

As in the previous case, the poles of the transmission coefficient yields the same set of QNFs found in the previous section. Also, we observe the same behavior described in the previous case for the reflection coefficient (105), transmission coefficient (106), and absorption cross section (107), i.e., we have found that the reflection coefficient is 1 at the low frequency limit, and for the high frequency limit this coefficient is null, with the behavior of the transmission coefficient being opposite with $R + T = 1$ (see Fig. (6)). Also, the absorption cross section is null in the low and high frequency limits, but there is a range of frequencies for which the absorption cross section is not null, and it also has a maximum value in the low frequency limit (see Fig. (7)). Furthermore, we observe that the absorption cross section can take higher values when the mass of the scalar field decreases (Fig. (7)) in the low frequency limit. However, beyond a certain value of the frequency the absorption cross section does not depend on the mass of the scalar field. It is worth noting that the absorption cross section does not depend on the angular momentum of the scalar field being the same for every value of κ .

V. CONCLUSIONS

In this work we calculated the QNFs of scalar field perturbations for the four-dimensional asymptotically Lifshitz black hole in conformal gravity with a spherical topology and dynamical exponent $z = 0$, where the anisotropic scale invariance corresponds to a space-like scale invariance with no transformation of time, for some special cases that depend on Q , and by imposing suitable boundary conditions at spatial infinity. These scalar fields are considered as mere test fields, without backreaction over the spacetime itself. Therefore, it is not necessary for such fields to have the same symmetries as the background spacetime. However, if one considers the backreaction of the matter fields over the spacetime, the relation between the symmetries of the spacetime and the matter fields is not trivial. For conformal gravity the gravitational field equations imply that the trace of the stress-energy tensor must vanish due to the Bach tensor is traceless.

Firstly, we analyzed massive scalar field perturbations minimally coupled to curvature, which does not have the same symmetries as the background spacetime due to the trace of stress-energy tensor is not null. The first case studied corresponds to a scalar field without angular momentum ($\kappa = 0$), and we found that there is a spectrum of quasinormal frequencies for which the scalar field becomes null at spatial infinity. These frequencies are purely




FIG. 6: The reflection coefficient R (solid curve), the transmission coefficient T (dashed curve), $R + T$ (thick curve) and the absorption cross section σ_{abs} (dotted curve) as a function of ω , for $m = 1$, $\ell = 1$ and $\kappa = 0$.




FIG. 7: The behavior of σ_{abs} as a function of ω , for $\kappa = 0$, $\ell = 1$ and $m = 1, 1.5, 2$.

imaginary and negative for $m = 0$; however, for $m \neq 0$ some QNFs are imaginary and positive. Another case we analyzed corresponds to $Q = \pm\infty$, where we found a spectrum of QNFs that respect the Dirichlet boundary condition; however, some of them have a positive imaginary part. Therefore, the black hole is unstable under massive scalar field perturbations and stable under massless scalar field perturbations. Also, we analyzed the extremal case $Q = 1$, for which we found that there are no QNMs as in [63], where the authors demonstrated the absence of QNMs in the extremal BTZ black hole. However, it was shown that it is possible to construct the QNMs of three-dimensional extremal black holes algebraically as the descendants of the highest weight modes [64], with hidden conformal symmetry being an intrinsic property of the extremal black hole. Also, it is worth mentioning that QNMs for extremal black holes are not always absent, for instance see [65], where the authors reported the presence of QNMs for extremal BTZ black holes in topologically massive gravity.

On the other hand, because the gravitational field equations imply that the trace of the stress-energy tensor must vanish, we also considered scalar field perturbations conformally coupled to curvature which have a traceless stress-energy tensor, and we showed that the imaginary part of the QNFs calculated is negative, what guaranties the stability of the Lifshitz black hole under conformally coupled scalar field perturbations, this was shown by using the improved AIM and analytical solutions. This behavior is similar to the studied in [66] for a three-dimensional Lifshitz black hole in conformal gravity.

Finally, we focused our attention to the minimally coupled scalar field case and we computed the reflection and transmission coefficients and the absorption cross section, and we showed numerically that the absorption cross section vanishes at the low and high frequency limits. Therefore, a wave emitted from the horizon, with low or high frequency, does not reach the spatial infinity and is totally reflected, because the fraction of particles penetrating the potential barrier vanishes. However, we have shown there is a range of frequencies where the absorption cross section is not null. The reflection coefficient is 1 at the low frequency limit and for the high frequency limit this coefficient is null, with the behavior of the transmission coefficient being opposite with $R + T = 1$. Also, we have shown that the absorption cross section increases if the mass of the scalar field decreases in the low frequency limit; however, beyond a certain value of the frequency the absorption cross section does not depend on the mass of the scalar field. Furthermore, we have shown that the absorption cross section does not depend on the angular momentum.

Acknowledgments

The authors would like to thank the referee for useful comments and constructive suggestions. This work was funded by the Comisión Nacional de Investigación Científica y Tecnológica through FONDECYT Grants 11140674 (PAG), 11121148 (YV, MC) and also by the Dirección de Investigación, Universidad de La Frontera (EC, MC). P. A. G. acknowledges the hospitality of the Universidad de La Serena where part of this work was undertaken. P. A. G. and Y. V. acknowledge the hospitality of the National Technical University of Athens.

-
- [1] S. Kachru, X. Liu and M. Mulligan, *Phys. Rev. D* **78**, 106005 (2008) [arXiv:0808.1725 [hep-th]].
 - [2] S. A. Hartnoll, J. Polchinski, E. Silverstein and D. Tong, *JHEP* **1004**, 120 (2010) [arXiv:0912.1061 [hep-th]].
 - [3] E. J. Brynjolfsson, U. H. Danielsson, L. Thorlacius and T. Zingg, *J. Phys. A* **43**, 065401 (2010) [arXiv:0908.2611 [hep-th]].
 - [4] S. J. Sin, S. S. Xu and Y. Zhou, *Int. J. Mod. Phys. A* **26**, 4617 (2011) [arXiv:0909.4857 [hep-th]].
 - [5] F. A. Schaposnik and G. Tallarita, *Phys. Lett. B* **720**, 393 (2013) [arXiv:1210.8358 [hep-th]].
 - [6] D. Momeni, R. Myrzakulov, L. Sebastiani and M. R. Setare, arXiv:1210.7965 [hep-th].
 - [7] Y. Bu, *Phys. Rev. D* **86**, 046007 (2012) [arXiv:1211.0037 [hep-th]].
 - [8] V. Keranen and L. Thorlacius, *Class. Quant. Grav.* **29**, 194009 (2012) [arXiv:1204.0360 [hep-th]].
 - [9] Z. Zhao, Q. Pan and J. Jing, *Phys. Lett. B* **735**, 438 (2014) [arXiv:1311.6260 [hep-th]].
 - [10] J. W. Lu, Y. B. Wu, P. Qian, Y. Y. Zhao and X. Zhang, *Nucl. Phys. B* **887**, 112 (2014) [arXiv:1311.2699 [hep-th]].
 - [11] G. Tallarita, *Phys. Rev. D* **89**, no. 10, 106005 (2014) [arXiv:1402.4691 [hep-th]].
 - [12] G. T. Horowitz and B. Way, *Phys. Rev. D* **85** (2012) 046008 [arXiv:1111.1243 [hep-th]].
 - [13] D. O. Devecioglu and O. Sarioglu, *Phys. Rev. D* **83** (2011) 021503 [arXiv:1010.1711 [hep-th]].
 - [14] D. O. Devecioglu and O. Sarioglu, *Phys. Rev. D* **83** (2011) 124041 [arXiv:1103.1993 [hep-th]].
 - [15] H. A. Gonzalez, D. Tempo and R. Troncoso, *JHEP* **1111** (2011) 066 [arXiv:1107.3647 [hep-th]].
 - [16] Y. S. Myung and T. Moon, *Phys. Rev. D* **86** (2012) 024006 [arXiv:1204.2116 [hep-th]].
 - [17] Y. S. Myung, *Eur. Phys. J. C* **72** (2012) 2116 [arXiv:1203.1367 [hep-th]].
 - [18] E. Ayon-Beato, A. Garbarz, G. Giribet and M. Hassaine, *Phys. Rev. D* **80** (2009) 104029 [arXiv:0909.1347 [hep-th]].
 - [19] R. -G. Cai, Y. Liu and Y. -W. Sun, *JHEP* **0910** (2009) 080 [arXiv:0909.2807 [hep-th]].
 - [20] E. Ayon-Beato, A. Garbarz, G. Giribet and M. Hassaine, *JHEP* **1004** (2010) 030 [arXiv:1001.2361 [hep-th]].
 - [21] M. H. Dehghani and R. B. Mann, *JHEP* **1007** (2010) 019 [arXiv:1004.4397 [hep-th]].
 - [22] R. B. Mann, *JHEP* **0906** (2009) 075 [arXiv:0905.1136 [hep-th]].
 - [23] K. Balasubramanian and J. McGreevy, *Phys. Rev. D* **80** (2009) 104039 [arXiv:0909.0263 [hep-th]].
 - [24] G. Bertoldi, B. A. Burrington and A. Peet, *Phys. Rev. D* **80** (2009) 126003 [arXiv:0905.3183 [hep-th]].
 - [25] M. Bravo-Gaete and M. Hassaine, *Phys. Rev. D* **89** (2014) 10, 104028 [arXiv:1312.7736 [hep-th]].
 - [26] A. Alvarez, E. Ayon-Beato, H. A. Gonzalez and M. Hassaine, *JHEP* **1406** (2014) 041 [arXiv:1403.5985 [gr-qc]].
 - [27] K. S. Stelle, *Phys. Rev. D* **16** (1977) 953.
 - [28] K. S. Stelle, *Gen. Rel. Grav.* **9** (1978) 353.
 - [29] P. D. Mannheim, *Found. Phys.* **37** (2007) 532 [hep-th/0608154].
 - [30] P. A. M. Dirac, "Lectures on Quantum Mechanics", Belfer Graduate School of Science, Yeshiva University Press, New York (1964).
 - [31] A. Pais and G. E. Uhlenbeck, *Phys. Rev.* **79** (1950) 145.
 - [32] T. Regge and J. A. Wheeler, *Phys. Rev.* **108**, 1063 (1957).
 - [33] F. J. Zerilli, *Phys. Rev. D* **2**, 2141 (1970).
 - [34] F. J. Zerilli, *Phys. Rev. Lett.* **24**, 737 (1970).
 - [35] K. D. Kokkotas and B. G. Schmidt, *Living Rev. Rel.* **2**, 2 (1999) [gr-qc/9909058].
 - [36] H. -P. Nollert, *Class. Quant. Grav.* **16**, R159 (1999).
 - [37] R. A. Konoplya and A. Zhidenko, *Rev. Mod. Phys.* **83**, 793 (2011) [arXiv:1102.4014 [gr-qc]].
 - [38] J. M. Maldacena, *Adv. Theor. Math. Phys.* **2**, 231 (1998) [hep-th/9711200].
 - [39] D. Birmingham, I. Sachs and S. N. Solodukhin, *Phys. Rev. Lett.* **88**, 151301 (2002) [hep-th/0112055].
 - [40] M. Alishahiha, M. R. Mohammadi Mozaffar and A. Mollabashi, *Phys. Rev. D* **86** (2012) 026002 [arXiv:1201.1764 [hep-th]].
 - [41] M. Catalan, E. Cisternas, P. A. Gonzalez and Y. Vazquez, *Eur. Phys. J. C* **74** (2014) 3, 2813 [arXiv:1312.6451 [gr-qc]].
 - [42] B. Cuadros-Melgar, J. de Oliveira and C. E. Pellicer, *Phys. Rev. D* **85**, 024014 (2012) [arXiv:1110.4856 [hep-th]].
 - [43] P. A. Gonzalez, J. Saavedra and Y. Vazquez, *Int. J. Mod. Phys. D* **21**, 1250054 (2012) [arXiv:1201.4521 [gr-qc]].
 - [44] P. A. Gonzalez, F. Moncada and Y. Vazquez, *Eur. Phys. J. C* **72**, 2255 (2012) [arXiv:1205.0582 [gr-qc]].
 - [45] R. Becar, P. A. Gonzalez and Y. Vazquez, *Int. J. Mod. Phys. D* **22**, 1350007 (2013) [arXiv:1210.7561 [gr-qc]].
 - [46] A. Giacomini, G. Giribet, M. Leston, J. Oliva and S. Ray, *Phys. Rev. D* **85** (2012) 124001 [arXiv:1203.0582 [hep-th]].
 - [47] J. M. Maldacena and A. Strominger, *Phys. Rev. D* **55** (1997) 861 [hep-th/9609026].
 - [48] T. Harmark, J. Natario and R. Schiappa, *Adv. Theor. Math. Phys.* **14** (2010) 727 [arXiv:0708.0017 [hep-th]].
 - [49] S. Lepe, J. Lorca, F. Pena and Y. Vazquez, *Phys. Rev. D* **86** (2012) 066008 [arXiv:1205.4460 [hep-th]].
 - [50] M. Olivares, Y. Vazquez, J. R. Villanueva and F. Moncada, arXiv:1306.5285 [gr-qc].

- [51] M. Olivares, German Rojas, Y. Vasquez and J. R. Villanueva, *Astrophys. Space Sci.* **347** (2013) 83 [arXiv:1304.4297 [gr-qc]].
- [52] J. R. Villanueva and Y. Vasquez, *Eur. Phys. J. C* **73** (2013) 2587 [arXiv:1309.4417 [gr-qc]].
- [53] H. Lu, Y. Pang, C. N. Pope and J. F. Vazquez-Poritz, *Phys. Rev. D* **86** (2012) 044011 [arXiv:1204.1062 [hep-th]].
- [54] I. Smolić, arXiv:1501.04967 [gr-qc].
- [55] M. Abramowitz and A. Stegun, *Handbook of Mathematical functions*, (Dover publications, New York, 1970).
- [56] P. P. Fiziev, *J. Phys. A: Math. Theor.* **43** (2010) 08. [arXiv:0904.0245[math-ph]].
- [57] H. T. Cho, A. S. Cornell, J. Doukas and W. Naylor, *Class. Quant. Grav.* **27** (2010) 155004 [arXiv:0912.2740 [gr-qc]].
- [58] H. Ciftci, R. L. Hall, and N. Saad, *J. Phys. A* **36**(47), 11807-11816 (2003).
- [59] H. Ciftci, R. L. Hall and N. Saad, *Phys. Lett. A* **340** (2005) 388.
- [60] H. T. Cho, A. S. Cornell, J. Doukas, T. R. Huang and W. Naylor, *Adv. Math. Phys.* **2012** (2012) 281705 [arXiv:1111.5024 [gr-qc]].
- [61] C. Y. Zhang, S. J. Zhang and B. Wang, arXiv:1501.03260 [hep-th].
- [62] T. Barakat, *Int. J. Mod. Phys. A* **21** (2006) 4127.
- [63] J. Crisostomo, S. Lepe and J. Saavedra, *Class. Quant. Grav.* **21** (2004) 2801 [hep-th/0402048].
- [64] B. Chen and J. -j. Zhang, *Phys. Lett. B* **699** (2011) 204 [arXiv:1012.2219 [hep-th]].
- [65] H. R. Afshar, M. Alishahiha and A. E. Mosaffa, *JHEP* **1008** (2010) 081 [arXiv:1006.4468 [hep-th]].
- [66] M. Catalán and Y. Vásquez, *Phys. Rev. D* **90**, no. 10, 104002 (2014) [arXiv:1407.6394 [gr-qc]].